

OCCUPATIONAL STRESS AND BURNOUT PREVENTION: A MULTILEVEL FRAMEWORK FOR WORKPLACE WELLBEING

Bistra Dimitrova

Prof, D. Sc. Faculty of public health, health care and Tourism

National Sports Academy "Vassil Levski", Bulgaria

E-mail: dimitrova.bistra@yahoo.com / ORCID.ID: 0000-0003-3152-9831

Abstract

Occupational stress and burnout represent challenges to public health, workforce sustainability, and organizational performance. Public discussions often combine established psychological concepts with personal testimony, commercial wellness claims, and recommendations of uncertain validity. This article systematizes source material concerning stress, burnout, panic-like symptoms, and self-regulation, distinguishing conceptually supported propositions from claims requiring clinical verification. A qualitative thematic content analysis was conducted. Statements were coded by conceptual domain, analytical level, intervention type, and evidentiary status, while repetition, promotional language, and first-person testimony were excluded. Five interrelated themes emerged: the adaptive and maladaptive dimensions of stress; occupational mismatch and burnout; the multidimensional nature of panic-like responses; self-regulation and lifestyle practices; and the boundary between supportive wellness measures and clinical treatment. Stress was not identified as uniformly harmful; however, prolonged exposure combined with inadequate recovery, limited autonomy, weak social support, and conflict between personal and organizational values may contribute to exhaustion and impaired functioning. Breathing regulation, progressive muscle relaxation, physical activity, social connection, and work-life boundaries may support recovery, whereas unverified remedies cannot be considered established treatments. Effective management requires an integrated model combining individual skills, organizational prevention, and access to healthcare. Wellness interventions should complement evidence-based assessment and treatment rather than replace them.

Keywords: *occupational stress, panic symptoms, self-regulation, workplace wellness, mental health.*

Introduction

In today's fast-paced and highly competitive work environments, professional stress has become a pervasive phenomenon affecting individuals across industries and organisational levels (2, 4, 6). Professional stress, commonly referred to as occupational or work-related stress, arises when job demands exceed an individual's capacity to cope or adapt, leading to physical, emotional, and behavioral strain (1, 5). While a certain level of stress can serve as a motivator, chronic or excessive stress has far-reaching implications for both employees and organisations (12, 14). Stress is simultaneously a biological response, a psychological experience and a social phenomenon (17, 18). It emerges when environmental demands are appraised as challenging or threatening and when available resources appear insufficient for effective adaptation (15, 16). This broad definition is important because it avoids treating every stress response as pathological. Short-term activation may mobilise attention and action, whereas persistent activation without adequate recovery may undermine wellbeing and performance (10, 19). Classical adaptation model describes stress through alarm, resistance and exhaustion. Although contemporary research has refined this sequence, the model remains useful for differentiating temporary mobilization from prolonged strain. Situational stress tends to diminish after the stressor is removed (11). Chronic stress persists

across time or continues after the original demand has ended. Post-traumatic responses may arise after exposure to an overwhelming event and require a distinct clinical assessment rather than being treated as an extension of workplace tension (9). Occupational stress occurs when job demands are poorly aligned with the worker's abilities, resources, autonomy or needs. The imbalance may involve excessive workload, time pressure, emotional demands, ambiguous roles, insufficient support, limited control or disruption of the boundary between work and private life (21-23). Its consequences may be emotional, cognitive, behavioral and somatic. Reported manifestations include fatigue, dissatisfaction, irritability, sleep disturbance, concentration difficulties, reduced performance and maladaptive coping (3). Such manifestations are not specific to one disorder and should therefore be interpreted within the person's broader medical and psychosocial context (20).

Burnout is more specific than stress. It refers to a work-related pattern characterized principally by exhaustion, psychological distancing or cynicism, and reduced professional efficacy (22). The source text presents burnout as the final stage of prolonged stress; however, the relationship is better understood as conditional rather than automatic. Exposure to stress may increase vulnerability, but occupational context, recovery opportunities, personal resources and organizational practices influence whether burnout develops. A useful organizational perspective is provided by six areas of potential mismatch: workload, control, reward, community, fairness and values (8). These areas shift attention away from interpretations that locate responsibility exclusively within the individual. A worker may practice relaxation and maintain healthy routines yet remain exposed to structurally excessive demands or ethical conflict. Sustainable prevention therefore requires changes in work design as well as individual coping support. The supplied text also connects stress with panic-like experiences, hydration, meditation, physical activity, herbal preparations and folk remedies. These elements represent different categories of knowledge. Some are established self-regulation practices; some are plausible but context-dependent health behaviors; others are personal testimonies or commercial claims. Combining them without qualification risks giving all statements an equivalent scientific status (13). The purpose of the present conceptual article is therefore threefold: first, to organize the material into a coherent model of occupational stress and burnout; second, to identify supportive self-regulation strategies that can be framed responsibly; and third, to clarify the boundary between wellness promotion and clinical treatment. The analysis does not test treatment effectiveness. It evaluates how the source material can be transformed into a defensible framework and identifies propositions requiring empirical investigation.

Research questions

The analysis addressed three questions: (1) Which core concepts structure the source account of stress and burnout? (2) Which personal and organisational responses are proposed? (3) Which claims require qualification because they rely on anecdote, promotional language or insufficient evidence?

Methods

2.1 Study design

A qualitative, document-based thematic synthesis was conducted. The unit of analysis was a Bulgarian-language source text addressing stress, occupational burnout, panic-like experiences and strategies presented as supportive or therapeutic. The design was appropriate because the material did not contain a defined sample, standardized measurements, an intervention protocol or outcome data that would support quantitative analysis. Accordingly, the article reports conceptual findings rather than clinical efficacy estimates.

2.2 Analytical procedure

The text was reviewed iteratively. During initial coding, statements were assigned to descriptive categories: definition of stress; temporal pattern; adaptation process; occupational antecedent; burnout characteristic; emotional, physiological or behavioral manifestation; self-regulation strategy; lifestyle recommendation; natural-product claim; and personal testimony. During focused coding, related categories were consolidated into higher-order themes. Attention was given to causal language, generalization, duplication and the difference between symptom support and disease treatment. A second analytical layer classified each proposition by evidentiary status. Conceptually established statements were retained in neutral academic language. Plausible but context-sensitive recommendations were presented as supportive measures rather than universally effective interventions. Personal accounts were treated as illustrations of lived experience and not as proof. Brand-specific claims, prices, dosing instructions and assertions of cure were excluded because the source provided no verifiable study design or safety evaluation.

2.3 Scope and limitations

The analysis was limited to the supplied material and did not constitute a systematic review of external literature. No bibliographic verification, risk-of-bias assessment or meta-analysis was undertaken. The absence of primary data precludes conclusions about prevalence, causality or comparative effectiveness. The resulting themes should be interpreted as a structured conceptual model and as a basis for future empirical research. Because the source included health-related advice, editorial transformation also required attention to scientific responsibility. The manuscript does not diagnose readers, prescribe products or substitute for professional care. Serious or persistent anxiety, recurrent panic episodes, suicidal thinking, chest pain, fainting, severe breathlessness or marked functional deterioration require timely assessment by qualified health professionals.

Results

Four criteria guided the transformation of the material. First, claims had to be proportionate to the information available. Second, stress, burnout, panic attacks and psychiatric disorders

were not treated as interchangeable constructs. Third, non-pharmacological self-care was distinguished from clinical management. Fourth, recommendations with potential safety implications were either qualified or omitted. For example, breathing into a paper bag was not retained as a general recommendation because acute breathlessness or chest symptoms may have causes other than panic, and rebreathing may be inappropriate in some situations (tab.1).

Table 1. Recommendations with potential safety implications

Analytical dimension	Operational question	Treatment in the synthesis
Conceptual coherence	Does the statement distinguish stress, burnout and panic symptoms?	Retain after clarification.
Evidence status	Is it a general concept, plausible strategy, testimony or promotional claim?	Label, qualify or exclude accordingly.
Causal precision	Does the text claim that one factor directly causes or cures a condition?	Reduce to an association unless supported by data.
Clinical safety	Could literal application delay care or create harm?	Add a safety boundary or omit.
Level of intervention	Is responsibility individual, organizational or clinical?	Integrate within a multilevel model.

3.1 Stress as an adaptive process with cumulative costs

The first theme presents stress as neither intrinsically harmful nor fully avoidable. Acute activation can support survival and goal-directed behavior by increasing vigilance and mobilizing resources. Risk emerges when demands remain intense, unpredictable or uncontrollable and recovery is insufficient. The source distinguishes situational, chronic and post-event patterns, suggesting that duration and recovery are central analytical variables. This distinction supports a continuum model in which the same adaptive system may become burdensome when activation is repeated or prolonged.

3.2 Occupational mismatch and burnout

The second theme concerns mismatch between the individual and the organization. Six recurrent domains were extracted: demands relative to capacity; autonomy relative to control; effort relative to recognition; quality of workplace relationships; perceived fairness; and compatibility between personal ethics and job requirements. These domains converge on the proposition that burnout is not merely a failure of personal resilience. It may indicate a sustained disruption in the relationship between employee and work environment. The manifestations described in the source span emotional exhaustion, diminished motivation, cognitive difficulty, behavioral withdrawal and physical complaints. The breadth of this list

requires caution: many symptoms are nonspecific and may reflect medical conditions, anxiety, depression, sleep disturbance or other factors. The analytically robust result is therefore not that burnout directly causes every listed condition, but that prolonged occupational strain may coexist with multidimensional impairment and warrants assessment at both individual and organizational levels.

3.3 Panic-like responses as multidimensional events

The third theme groups panic-like experiences into emotional, physiological and behavioral components. Emotional features include intense fear and catastrophic thoughts; physiological features may include palpitations, dizziness, breathlessness, abdominal discomfort and muscular tension; behavioral responses may include avoidance and sleep disruption. This three-domain representation is useful for explaining why panic is experienced as a whole-body event. Nevertheless, symptom overlap with cardiopulmonary, endocrine, neurological and other conditions means that first or atypical episodes should not automatically be self-classified as panic (fig.1).




Figure 1. Integrated multilevel response to occupational stress and burnout.

Figure 1. Integrated multilevel response to occupational stress and burnout.

3.4 Self-regulation and recovery

The fourth theme comprises strategies intended to reduce arousal and restore a sense of control. The most defensible measures in the source include slow, comfortable breathing without prolonged breath-holding; progressive muscle relaxation; attention-grounding; regular physical activity; reduction of excessive stimulant intake; consistent sleep routines; constructive social support; and cultivation of interests outside work. Meditation may also serve as a structured attentional practice when adapted to the individual and, for beginners or vulnerable participants, introduced with appropriate guidance.

3.5 Boundary between wellness support and treatment

The fifth theme emerged from tension between low-risk wellbeing practices and claims that foods, herbs or branded supplements treat panic, depression or severe psychiatric conditions. The source includes testimonials and product descriptions but no controlled evidence, diagnostic verification, interaction assessment or systematic adverse-event reporting. Consequently, these materials cannot support efficacy conclusions. Natural origin does not establish safety, and herbal ingredients may interact with medication, affect pregnancy or lactation, and be unsuitable for particular medical conditions (tab. 2).

Table 2. Differential assessment of levels and objectives with examples

Level	Primary objective	Examples derived from the synthesis
Individual	Improve regulation and recovery.	Sleep routine, physical activity, relaxation, social connection, work-life boundaries.
Organizational	Reduce preventable psychosocial hazards.	Workload review, autonomy, recognition, fairness, supportive teams, ethical alignment.
Clinical	Assess persistent, severe or ambiguous symptoms.	Differential assessment, evidence-based psychotherapy, medical review and monitored treatment.

3.6 Implications for research and practice

Future studies should operationalize the proposed model using validated measures of perceived stress, burnout, work characteristics, sleep, recovery and functioning. Prospective designs could examine whether mismatch domains predict changes in burnout and whether organizational improvements modify this relationship. Intervention trials should specify the population, comparator, duration, adherence, adverse events and clinically meaningful outcomes. Natural products require standardized composition, interaction screening and independent safety monitoring.

For practice, the findings support stepped care. Universal prevention may include healthy work design and basic recovery education. Selective support may target teams or employees exposed to sustained demands. Indicated care should provide prompt access to qualified psychological or medical services. Evaluation should measure not only symptom change but also workload, absenteeism, retention, psychological safety and work quality.

Discussion

The thematic synthesis supports a multilevel interpretation of stress-related problems. At the individual level, stress involves appraisal, physiological activation and coping behavior. At the occupational level, workload, control, reward, social climate, fairness and values shape exposure and recovery. At the clinical level, persistent anxiety, panic symptoms, depressive features and physical complaints may require formal assessment. Treating these levels as interchangeable can produce two opposite errors: medicalizing ordinary adaptive strain or

minimizing clinically significant distress as a lifestyle problem. The strongest conceptual contribution of the source is its recognition that stress may be protective in the short term but damaging when prolonged. This prevents the unrealistic objective of eliminating all stress. A more appropriate goal is regulatory flexibility: the capacity to mobilize when necessary and return to baseline after the demand passes. Recovery should therefore be viewed as an active component of performance rather than as inactivity or lack of commitment. The six mismatch domains offer a practical bridge between theory and organizational policy. A wellness program focused only on breathing exercises, meditation or supplements may have limited impact if employees remain exposed to chronic overload, conflicting expectations or unfair treatment. Conversely, organizational redesign without accessible individual support may overlook differences in vulnerability, coping resources and life circumstances. Integrated prevention requires both approaches and should include mechanisms for early identification, confidential referral and follow-up. The source's treatment of panic-like experiences illustrates the need for careful communication. Grounding, gentle breathing and muscle relaxation can help some individuals regulate acute anxiety. However, categorical assurances that every episode is harmless are inappropriate because similar symptoms may occur in medical emergencies. Public-facing wellness materials should advise individuals to seek urgent evaluation for new, severe or unusual chest pain, loss of consciousness, marked breathing difficulty, neurological symptoms or other alarming features. Recurrent episodes also merit professional assessment because effective psychological and medical options are available. A second concern is the transformation of personal testimony into therapeutic recommendation. Lived experience can generate hypotheses and illuminate patient perspectives, but it cannot establish efficacy, dosage or safety. The presence of a named product, consumer price or anecdotal recovery narrative introduces promotional bias. For a scholarly publication, such content should either be studied explicitly as health communication or excluded from treatment recommendations. The same principle applies to broad claims that a natural preparation can treat heterogeneous conditions ranging from stress to severe mental illness. Hydration deserves similar qualification. Adequate fluid intake contributes to normal physiological functioning and may support wellbeing, particularly during heat, exercise or illness. It should not, however, be presented as a primary causal explanation or cure for cellular aging, burnout, panic disorder or depression. Reductionist explanations may be attractive because they suggest a simple solution, but they obscure the biological, psychological, occupational and social determinants involved.

Conclusion

Stress is an unavoidable component of adaptation, yet its consequences depend on intensity, duration, appraisal, available resources and opportunities for recovery. The supplied material contains an important central insight: prolonged imbalance between demands and resources can affect emotional wellbeing, cognitive functioning, behavior and occupational

performance. The analysis further indicates that burnout is best understood as a work-related phenomenon situated at the intersection of the person and the organizational environment. Five themes organize the proposed framework: adaptive versus cumulative stress; person–work mismatch; multidimensional panic-like responses; self-regulation and recovery; and the distinction between wellness support and clinical treatment. Together they show that effective prevention cannot rely on a single technique. Individual skills such as relaxation, physical activity, sleep regulation and social connection may support recovery, but they do not compensate indefinitely for excessive workload, low autonomy, lack of recognition, unfairness or value conflict.

The source also demonstrates why academic health communication must evaluate evidentiary status. Personal narratives can be meaningful, and traditional practices may merit investigation, but neither testimonials nor commercial descriptions establish therapeutic effectiveness. Claims about herbal preparations, foods or supplements should not be generalized to the treatment of panic disorder, depression or severe psychiatric conditions without appropriate clinical evidence. Product use additionally requires attention to contraindications, interactions and vulnerable populations.

A responsible workplace-wellness model should therefore include three coordinated components. The first is organizational prevention through realistic demands, meaningful control, fair recognition, supportive relationships and ethical coherence. The second is individual capacity-building through accessible education in recovery, emotional regulation and help-seeking. The third is a clear clinical pathway for persistent, severe or diagnostically uncertain symptoms. Each component addresses a different source of risk, and none is sufficient in isolation.

This conceptual synthesis does not provide prevalence estimates or treatment-effect sizes. Its contribution is methodological and practical: it transforms heterogeneous source material into a structured model while preventing anecdotal and promotional claims from acquiring an unwarranted scientific status. The model can inform the development of questionnaires, workplace interventions and prospective studies, but its propositions require empirical testing in defined occupational populations.

In conclusion, the most sustainable response to stress and burnout is neither the elimination of all challenges nor the transfer of responsibility entirely to the worker. It is the creation of environments in which demands are proportionate, recovery is possible, support is accessible and clinical concerns are recognized early. Within such an environment, wellness practices can serve their appropriate purpose: supporting resilience and quality of life while complementing evidence-based occupational and healthcare interventions.

Acknowledgement:

Conflict of Interest: The author and the institution declared no conflict of interest.

Financial Disclosure: The article is under the implementation of project BG16RFPR002-1.014-0011 Sustainable Development of the Center of Excellence "Heritage BG," funded under the grant aid procedure BG16RFPR002-1.014 "Sustainable Development of Centres of Excellence and Centres of

Competence, Including Specific Infrastructures or Their Associations from the National Roadmap for Scientific Infrastructure”.

Declarations of informed consent have been signed regarding the publication of survey data. ChatGPT is used for English grammar revision and translation.

References

1. Aleksandrova, V. (2021a). *Adaptirano pluvane za detsa s uvrezhdaniya* [Adapted swimming for children with disabilities] [Monograph, in Bulgarian]. IPRESS.
2. Aleksandrova, V. (2021b). The effect of swimming on health and physical activity in children with visual disabilities. *Journal of Environmental Protection and Ecology*, 22(2), 847–857.
3. Aleksandrova, V. (2022). Wellness aqua programs through adapted physical activity and sports for children with special needs. *Research Announcements Heritage BG*, 3, 82–89.
4. Angelcheva, M. (2019). Effects of combination between relax massage and Dr Bach flower remedies on everyday stress. In *Proceedings of the International Scientific Congress "Applied Sports Sciences" & Balkan Scientific Congress "Physical Education, Sports, Health"* (p.348–352). NSA Press.
5. Angelcheva, M., & Petkova, S. (2019). Effects of combination between relax massage and Dr Bach flower remedies on everyday stress. In *Proceedings of the International Scientific Congress "Applied Sports Sciences" and Balkan Scientific Congress "Physical Education, Sports, Health"* (pp. 348–352). NSA Press.
6. Chipeva, M. (2018). *Aerobika: Istoriya, izsledvaniya, obuchenie* [Aerobics: History, research and education] [In Bulgarian]. NSA Press.
7. Chipeva, M. (2018a). *Aerobno byagane* [Aerobic running] [In Bulgarian]. NSA Press.
8. Chipeva, M. (2019). *Kango jumps aerobika* [In Bulgarian]. Avangard Prima.
9. Dimitrova, N. (2017). The factor motivation for judo training with children. *Research in Kinesiology*, 45(2), 161–163.
10. Dimitrova, N. (2019). Resistance while managing its own inertial power field. *Trakia Journal of Sciences*, 17(1), 631–634.
11. Dimitrova, N. (2019a). Reactions of the motor system in local and global activities. *Trakia Journal of Sciences*, 17(1), 635–637.
12. Dimitrova, N. (2020). Control of the spatial structure in preschool children. *Trakia Journal of Sciences*, 18(1), 854–856.
13. Dimitrova, N. (2025). Specificity of balance in disposal (weightlifting). *German International Journal of Modern Science*, 102, 63–66.
14. Dimitrova, N. (2026). Comparative biomechanical analysis of performance against a real target and a "kata" in karate. *Trakia Journal of Sciences*, 24(2). <https://doi.org/10.15547/tjs.2026.02.012>
15. Nesheva, I. (2022). Social benefits of wellness motor activities for women with normal pregnancy. *Smart Innovations in Recreational, Wellness Industry and Niche Tourism*, 4(1–2), 22–27.
16. Nesheva, I. (2023). Pregnancy and wellbeing. *Trakia Journal of Sciences*, 21(Supl. 1), 231–235. <https://doi.org/10.15547/tjs.2023.s.01.039>
17. Nesheva, I. (2023a). A wellness tool to perform healthy lifestyle practices in the Bulgarian school system. *Strategies for Policy in Science and Education*, 31(5), 560–572. <https://doi.org/10.53656/str2023-5-7-wel>

18. Polimenov, M. (2014). Innovation in SPA and enology as an opportunity to raise the competitiveness of tourist services. In *Proceedings of the International Tourist Forum "SPA and Wine"* (p. 163). South-West University "Neofit Rilski".
19. Polimenov, M. (2018). Technological innovations in niche tourism. *Smart Innovations in Recreational, Wellness Industry and Niche Tourism*, Special Issue, 31–35. <https://scjournal.globalwaterhealth.org/>
20. Polimenov, M. (2019). Transfer of innovation in service technology for increasing restaurant product quality. *Smart Innovations in Recreational, Wellness Industry and Niche Tourism*, 1(2), 29–35. <https://scjournal.globalwaterhealth.org/>
21. Tomova, T., & Angelov, B. (2023). Wellness practices for healthy ageing. *Trakia Journal of Sciences*, 21(Suppl. 1), 445–450. <https://doi.org/10.15547/tjs.2023.s.01.074>
22. Tomova, T. (2024). Modeling rituals through combined wellness influences. *Smart Innovations in Recreational, Wellness Industry and Niche Tourism*, 6(2), 15–24.
23. Tomova, T. (2025). The impact of manual wellness therapies and body massage: A stress relief effect. *German International Journal of Modern Science*, 105, 23–27.